

Urology Update from Jackson Hole

*Highlights from the 3rd Annual Jackson Hole Summer Urologic Conference
July 28–August 3, 2001, Jackson Hole, WY*

[*Rev Urol.* 2002;4(2):78–86]

© 2002 MedReviews, LLC

Key words: Anti-androgen monotherapy • Incontinence • Infertility treatments, cost/benefit • Prostate cancer • Prostatectomy • Ureteral stones • Urinary diversion

The 3rd annual Summer Jackson Hole Urologic Conference in Jackson Hole, WY once again combined an outstanding educational format with a spectacular venue. Conceived over two decades ago by Ralph Hopkins, a practicing urologist from Riverton, WY, the Jackson Hole meeting combines an internationally recognized faculty with an equally renowned critique panel. The faculty present a series of papers in their areas of expertise and the critique panel is afforded sufficient time to critically analyze and evaluate the presentations. This results in some good-natured ribbing but also allows critical assessment of the data presented.

Reviewed by Michael K. Brawer, MD,
Northwest Prostate Institute, Seattle, WA

Prostate Cancer

TURP-Detected Prostate Cancer

Gerald Andriole presented a number of interesting papers, primarily on carcinoma of the prostate, based on his vast experience at Washington University School of Medicine in St. Louis. In a paper on the frequency and histopathologic features of transurethral resection of the prostate (TURP)-detected prostate cancer, he first reviewed the literature on the incidence of transition zone carcinoma (see Table 1).^{1–7} He went on to describe his experience in the prospective evaluation of 259 men with lower urinary tract symptoms (LUTS) who underwent TURP. All men with either an abnormality on digital rectal examination or elevated prostate-specific antigen (PSA) underwent at least one set of peripheral

and/or transition zone biopsies that were negative for cancer. The incidence of cancer and the number of prior negative biopsies are shown in Table 2. These data demonstrate that a considerable portion of patients who have negative peripheral and indeed transition zone biopsy may harbor significant carcinoma of the prostate at the time of TURP. The decrease in incidence of TURP as we move more and more to pharmacologic and alternative therapies for LUTS owing to presumed benign prostatic hyperplasia (BPH) underscores the need to periodically evaluate those patients for prostate cancer.

Gleason Score 3+4 Versus 4+3

Dr. Andriole reviewed the differences between Gleason 3+4 and Gleason 4+3 in another presentation. He eval-

Table 1
Percent Prostate Cancer in Transition Zone Biopsies
if Prior Negative Prostate Biopsy

Repeat Positive Biopsy Rate	Prostate Cancer Only in Transition Zone	Author
37%	—	Lui et al ¹
31%	10%	Liu et al ²
32%	15%	Babaian ³
24%	3%	Bazimet et al ⁴
16%	10%	Keetch et al ⁵
21%	9%	Morote et al ⁶
18%	12%	Onder et al ⁷

uated 133 men with Gleason 3+4 and 24 men with Gleason 4+3 who were followed for a mean time of 38.6 months. Those men with Gleason 4+3 had the higher pathologic stage and larger tumor volume, although these differences were not statistically significant. The incidence of PSA recurrence for those men with Gleason 4+3 was significantly greater than those with Gleason 3+4 ($P = .0008$). These data confirm the observation from the Stanford group and others that increasing percentage of Gleason grade 4 and/or 5 connotes an adverse prognostic indication.

Radical Prostatectomy

In a paper addressing the use of selective cavernous nerve resection and sural nerve grafting for men undergoing radical prostatectomy, Dr. Andriole gave a balanced presentation. He noted that, although technically feasible, the indication for wide resection of the nerve vascular bundles is at best obscure. Using preoperative parameters may result in inappropriate nerve resection in 15% to 60% of cases. He concluded that until more is known about the appropriate factors to guide the decision for nerve resection, which hopefully will come from advances in staging

methodology, this procedure should be considered investigational.

The finding of positive surgical margins at the time of radical prostatectomy continues to occur all too often, and the appropriate intervention remains obscure. Dr. Andriole identified 33 men with isolated positive apical margins and compared them to a cohort of patients matched by age and preoperative PSA and biopsy Gleason parameters who were shown to have pathological T2 disease. Patients were evaluated for PSA failure at 1-, 3-, and 5-year intervals. PSA recurrence for survival at 5 years was 75% and 77%, respectively, for those with positive and negative api-

cal margins ($P > .10$). Dr. Andriole concluded that when controlling for other prognostic parameters a positive margin at the apex is not associated with untoward prognosis.

Percent of Positive Biopsies as a Predictor of Outcome

Gary Grossfeld, Assistant Professor of Urology at the University of California at San Francisco, also focused primarily on carcinoma of the prostate. One paper addressed the use of the percent of biopsies with cancer to predict tumor extent and outcome following radical prostatectomy. He used the Cancer of the Prostate Strategic Urologic Research Endeavor (CaPSURE) registry of patients throughout the United States. Univariate and multivariate analysis demonstrated that PSA at diagnosis, biopsy Gleason score, and the percentage of positive biopsies (number of positive biopsies/total number of biopsies) all were significant predictors of both extracapsular disease and disease recurrence. The percent of biopsies that revealed carcinoma was the most important predictor for the likelihood of extra prostatic disease and disease recurrence for those patients with intermediate or high-risk disease characteristics, based on clinical tumor stage, serum PSA, and

Table 2
TURP-Detected Prostate Cancer

	No. Patients	No. (%) with Prostate Cancer
No prior biopsy	40	6 (15)
One or more negative peripheral zone biopsy only	105	13 (13)
1 or more negative peripheral + T2 biopsy	144	11 (8)

Data from Ornstein et al.²⁶

TURP, transurethral resection of the prostate.

Table 3
Trials Comparing Bicalutamide Monotherapy with Castration

Author	N	Dose	Stage of Disease	Comparison	Result
Chodak et al ⁹	486	50 mg	Stage D2	Castration	Longer progression-free with castration
Bales and Chodak ¹⁰	1037	50 mg	Stage D2	Castration	Longer median survival with castration
Tyrrell et al ¹¹	1453	150 mg	Metastatic and locally advanced	Castration	Longer survival in M1 patients with castration
Boccardo et al ¹²	229	150 mg	Metastatic and locally advanced	Combined androgen blockade	No difference in progression-free or overall survival
Iversen et al ⁸	480	150 mg	Locally advanced	Castration	No difference in survival

biopsy grade. In a subset of low-risk patients, disease recurrence at 5 years was 15% for those with one third or less of the biopsies positive, as compared to 31% for those that had more positive biopsies. Similar findings were noted with intermediate- and higher-risk disease. The simplicity of simply identifying the number of biopsy cores that are positive renders this approach widely applicable, although certainly not as reproducible as a more sophisticated measurement, such as percent of cancer within a biopsy or millimeters of biopsies that showed carcinomas. Using percent of positive biopsies as a measure obviously has more general appeal. Obviously one confounder is the number of biopsies obtained, and this needs to be standardized for this method to be useful.

Intermittent Androgen Deprivation

Dr. Grossfeld provided an update on the University of California experience with intermittent androgen deprivation. Sixty-one patients were treated, including 34 patients with clinically localized disease and no prior treatment and 27 patients who had recurrent disease following local

therapy. In their study, the University of California group treats patients with 6–9 months of androgen deprivation during the initial cycle. They continue this for 1 to 2 months after PSA either reaches a nadir or becomes undetectable. Patients who achieved a PSA nadir less than 0.1 ng/mL (for those with prior definitive local therapy) or less than 4.0 ng/mL (for those with no prior local therapy) are eligible for going off therapy. Androgen deprivation is then reinstituted when serum PSA reaches a predetermined level either greater than 50% of the pretreatment level or greater than 10.0 ng/mL, or by patient request. With follow-up between 7 and 60 months following the start of treatment, with a mean of 30 months, patients received one to five treatment cycles with a median of 2 cycles and the median cycle length being 14 months. Importantly, patients were off androgen deprivation 45% of the time, although this time interval decreased with the increasing number of cycles. Among a small subset of patients in which quality of life was measured with a variety of instruments, the researchers noted clinically significant improvement in

health-related quality of life during the off-treatment episodes in the domains of vitality and fatigue, sexual function, and sexual bother.

Anti-Androgen Monotherapy

Dr. Grossfeld reviewed the experience with anti-androgen monotherapy in prostate cancer. A number of studies have reported on the use of bicalutamide in either the 50 or 150 mg per day dosing. These demonstrated that castration was superior irrespective of bicalutamide dose. Iversen and associates⁸ observed that in 480 patients with locally advanced non-metastatic prostate cancer and a median follow-up of 6.3 years, high-dose bicalutamide was reported to be equivalent to castration in terms of time-to-disease progression and overall survival. Significant benefit to the patients randomized to bicalutamide included improvement in sexual interest and physical capacity. Adverse side effects included hot flashes in the castration group and breast pain and gynecomastia in the bicalutamide group. Table 3 depicts several trials in which bicalutamide is compared with surgical or pharmacologic castration.^{8–12}

Management of Disease Following Radical Prostatectomy

The management of the man with positive surgical markers following radical prostatectomy is, as noted above, a confusing one. No definitive trials have been reported in which patients were properly randomized. Options include delayed radiation or androgen deprivation at the time of disease progression, immediate adjuvant radiation, or androgen deprivation. Each has its proponents. Grossfeld noted that although immediate radiation therapy in an adjuvant setting is certainly advocated by many, if not most, radiation oncologists, data from the University of Colorado and San Francisco demonstrate that if therapeutic radiation is given with a PSA below 2.0 ng/mL, the results are indistinguishable from patients receiving adjuvant radiation before PSA becomes detectable.¹³ Dr. Grossfeld described a decision analysis model to determine the preferred management of positive surgical margins.¹⁴ Using the average probability assessments from the literature, the

decision model recommended initial surveillance. Sensitivity analyses demonstrated the model was dependent on the probability of disease recurrence in patients followed expectantly as well as the efficacy of therapeutic radiation.

Urinary Diversion

Dr. Grossfeld gave an interesting talk on orthotopic urinary diversion in women requiring cystectomy. He noted that increase in experience demonstrated that this method is a safe and reliable method of urinary diversion in properly selected patients. Examination must include assurance that urethra is not involved in neoplasia. Stein and associates¹⁵ noted that the only risk factors for urethral involvement, based on 67 female cystectomy specimens, are tumor involvement of the bladder neck or the anterior vaginal wall. Dr. Grossfeld's summary of the literature may be found in Table 4.

Incontinence

Niall Galloway, Associate Professor

and Director of the Emory Continence Center in Atlanta, presented a number of papers on pelvic anatomy and the mechanisms of cystocele, rectocele, and urinary incontinence. In an illuminating presentation on the evaluation of the incontinent patient, Dr. Galloway cautioned that although traditional medical specialties have divided the organs of the pelvis into the gastrointestinal, gynecologic, and genital urinary "camps," that in fact these organs work together in health and often fail together in disease. He underscored the need for a broader approach to optimum management of patients with pelvic disorders. In addition to the evaluation of the patient, Dr. Galloway emphasized the importance of a full clinical history, a thorough physical examination, measurement of uroflow and post-void residual volume, and urine analysis. He has his patients keep a bladder diary before their visit.

One point that was illuminating to most of the audience was that the foot is innervated by S2 and S3, and thus examination of the feet gives an

Table 4
Continence Results for Women Undergoing Orthotopic Urinary Diversion Following Cystectomy

Author	Year	N	% Continent	% Catheterize	% Additional Treatment for Incontinence
Stein (personal communication)	2001	88	75%	34%	14%
Blute and Gburek ²⁷	1998	5	80%–100%	20%	—
Cancrini et al ²⁸	1995	7	71%–100%	—	—
Hautman et al ²⁹	1996	18	85%	70%	—
Jarolim et al ³⁰	1997	12	92%	8%	—
Mills and Studer ³¹	2000	15	80%–100%	7%	—
Shimogaki et al ³²	1999	8	88%	50%	—
Stenzl et al ³³	1997	20	79%–88%	4%	—
UCSF	2001	25	76%	8%	16%

UCSF data from Gary D. Grossfeld, MD.

Table 5
Spontaneous Passage Rates for Ureteral Stones

Size	Rate	Location	Rate
<4 mm	57%	Proximal	12%
4–6 mm	35%	Middle	22%
>6 mm	8%	Distal	45%

Data from Hubner et al.¹⁶

indication of the neuronal health or lack thereof of the pelvic floor and sphincters, which are innervated by S3, S4, and S5. Another physical finding that may demonstrate underlying pathology is asymmetry of the buttocks and gluteal folds in the standing position. Two-point discrimination, he feels, should be evaluated in the peri-anal and post-anal dermatomes and muscle tone and group of the circum-vaginal and anal sphincter muscles. The perineum should be examined with a full bladder in the standing as well as the supine position. The bladder scan provides an easy method of evaluating a post-void residual volume, which can be measured repeatedly.

Dr. Galloway advocates non-surgical treatments for all patients initially. These include fluid and bowel management, skin care, and containment issues. Careful drug history is important, and agents such as α -blockers should be avoided, as they result in incontinence. Dr. Galloway recommends a natural conservation management and evaluation. However, if the symptoms are moderate or severe, or if anatomical defects are present, urodynamic evaluation is important.

Endourology

Distal Ureteral Stones

Dr. Margaret Pearle, Associate Professor of Urology at the University of Texas Southwestern Medical Center, provided an update on a number of

endourologic advances. Perhaps no field of urology is today more controversial than the option of therapy for distal ureteral stones. She noted that observation is appropriate in select cases, and with ureteral stent placement to dilate the ureter, spontaneous passage is common. Ureteroscopy (URS) and shockwave lithotripsy (SWL) are both highly successful with low complication rates. Salvage therapy, either laparoscopic or open ureterolithotomy, are rarely needed. The rates for spontaneous passage of stones have been reviewed by Hubner et al¹⁶ (see Table 5). The use of a stent as definitive management is demonstrated by Leventhal et al.¹⁷ He randomized men to URS versus stent placement. Seventeen patients were initially managed with a ureteral stent. Eighty-three percent passed the stone at a mean of 6.6 days. There was no difference in stone size between groups. Dr. Pearle made the point that when SWL is considered for ureteral stone management, if a stent needs to be passed it may be just as easy to pass a 7 French ureteroscope and treat the stone definitively. Complications associated with SWL are primarily related to fragment passage, with sepsis occurring in 1% to 1.5%. No ureteral injury or stricture has been reported with SWL in the absence of manipulation or stem placement. Complications from URS include a perforation rate that is now less than 4% and late ureteral stricture formation in 2%.

Wollin and Denstedt¹⁸ randomized 58 patients with ureteral stones to receive a stent or not. At 1 week, flank and bladder symptoms were greater in patients with a stent. One hundred percent stone-free status was achieved in all patients, and one patient in each group required an emergency room visit or admission. These authors concluded that routine stent placement after uncomplicated URS for distal ureteral stones may be unnecessary. A number of studies have been reported in which patients were randomized to URS versus SWL for distal ureteral stones. A report by Deliveliotis et al¹⁹ randomized patients equally to URS versus SWL. For distal stones, stone-free rates were 100% in the URS group and 98% in the SWL group, with a 6% retreatment rate. SWL operating room time was 10 minutes shorter, and there was a 4% stricture rate in

Faculty, 3rd Annual Jackson Hole Summer Urologic Conference

Gerald L. Andriole, MD

Washington University,
School of Medicine
St. Louis, MO

Margaret S. Pearle, MD

Associate Professor of Urology
University of Texas Southwestern
Medical Center at Dallas
Dallas, TX

Gary D. Grossfeld, MD

Assistant Professor of Urology
University of California at San Francisco
San Francisco, CA

Niall T.M. Galloway, MD, FRCS

Associate Professor, Medical Director
The Emory Continence Center,
Department of Urology
Atlanta, GA

Lawrence S. Ross, MD

Clarence C. Saelhof Professor and Head,
Department of Urology
University of Illinois at Chicago
Chicago, IL

Table 6
Balloon Dilation: Review of the Literature

Author	N	Caliber Balloon	Size Stent (Fr)	Duration of Stent	Success	Follow-up
Johnson et al ³⁴	31	4–10 mm	5–14	≥2 d	58% (18/31)	≥6 mo
Smith ³⁵	5	—	—	—	39% (7/18)	—
Chang et al ²²	10	5–8 mm	8–16	4–8 wk	70% (7/10)	—
O'Brien et al ³⁶	14	4–6 mm	8.5	—	57% (8/14)	14 mo
Beckmann and Roth ³⁷	17	4–10 mm	7–10	4–8 wk	82% (14/17)	15 mo
Kramolowsky et al ³⁸	14	—	—	—	64% (9/14)	27 mo
Netto et al ³⁹	21	4–6 mm	8.5	Average 40 d	57% (12/21)	21 mo
Ravery et al ⁴⁰	11	10 mm	8	2.1 mo	40% (4/10)	16 mo
Total	123	—	—	—	59% (79/135)	—

the URS group, with none in the SWL group. Peschel et al²⁰ reported on 80 patients with distal ureteral stones that failed to pass in 3 weeks. They were randomized to SWL or URS. Stone-free rates were 90% in the SWL and 100% in the URS group. Operating room times were shorter for URS, as was time to becoming stone-free.

Ureteral Strictures

Dr. Pearle reviewed the endourologic management of ureteral strictures. She notes that URS is the most common cause today for these. Treatment options include balloon dilation, endoureterotomy, and ureteroneocystostomy. She reviewed the literature with respect to balloon dilation, as is shown in Table 6. Diagnostic factors associated with stricture duration have been reported by Beckmann²¹ for strictures of less than 3 months' duration. There was an 88% success rate, compared to 67% for those of longer status. Chang et al²² noted 91% success for strictures of less than 3 months versus 25% for those greater than 3 months. Chang²² also

noted the length of stricture would offer prognostic information, with 100% of stricture success with balloon dilation for those less than 1.5 cm versus 12.5% for those greater than 1.5 cm. Endoureterotomy technique was discussed. Dr. Pearle recommends a transmural endoscopic incision through the stricture extending into normal ureter proximally and distally. Stent placement for 6 weeks is required. A review of the literature

with respect to endoureterotomy may be found in Table 7. Complications from endoureterotomy include rare urinoma formation and ureteroenteric fistula. Vascular injuries also are rare but must be considered as a potential outcome. Dr. Pearle concluded that endourologic management of benign ureteral and ureteroenteric strictures provides reasonable first-line therapy. Endoureterotomy has a higher success rate than balloon dilation and the

Table 7
Endoureterotomy: Results

Author	N	Modality	Follow-up	Success
Schneider et al ⁴¹	12	Cold knife	15 mo	67% (8/12)
Meretyk et al ⁴²	13	Electrified cut	20 mo	62% (8/13)
Yamada et al ⁴³	20	Cold knife	18 mo	85% (17/20)
Cornud et al ⁴⁴	31	Electrified cut	>12 mo	71% (22/31)
Preminger et al ⁴⁵	40	Acucise	8.7 mo	53% (21/40)
Singal et al ⁴⁶	12	Ho:YAG	≥9 mo	75% (9/12)
Wolf et al ⁴⁷	38	Variety	28 mo	82% (31/38)
Total	166	—	—	70% (116/166)

latter is exceedingly prone to failure for ureteroenteric strictures. She underscored the need for prospective randomized trials to optimize treatment methods.

Lower Pole Ureteral Stones

Management of the lower pole ureteral stone remains controversial. Perhaps the most important study in this regard was the meta-analysis performed by Lingeman et al.²³ Thirteen shockwave studies and three percutaneous nephrolithotomies were evaluated. This study demonstrated that, controlling for stone size, patients with lower pole stones are 6.27 times more likely to be rendered stone-free with percutaneous approaches as compared to SWL. The




Figure 1. Algorithm for management of lower pole stones. URS, ureteroscopy; SWL, shockwave lithotripsy; PCNL, percutaneous nephrolithotomy.

Critique Panel, 3rd Annual Jackson Hole Summer Urologic Conference

Michael K. Brawer, MD

Director, Northwest Prostate Institute
Northwest Hospital
Seattle, WA

Elsbeth M. McDougall, MD

Professor of Urologic Surgery
Vanderbilt University Medical Center
Nashville, TN

Randall B. Meacham, MD

Associate Professor
Residency Program Director & Chief,
Division of Urology
Director of Male Reproductive
Medicine and Surgery
University of Colorado Health
Science Center
Denver, CO

Ian M. Thompson, MD

Professor and Chief
Division of Urology
University of Texas Health
Science Center
San Antonio, TX

Hunter Wessells, MD

Associate Professor
Department of Urology
University of Washington
Harbor View Medical Center
Seattle, WA

prospective randomized trial by Lingeman, in which 160 patients were randomized to SWL versus percutaneous nephrolithotomy (PCNL) and stratified with stone size, demonstrated that lithotripsy resulted in stone-free status in only 37%, compared to 95% of those having a percutaneous approach. Dr. Pearle's algorithm for the management of lower pole stones is shown in Figure 1. Dr. Pearle's conclusion as to the outcome and management of lower pole calculi is that for small stones (<10 mm), SWL is reasonable first-line therapy. For larger stones (>20 mm), PCNL is preferred. For stones intermediate in size (11–20 mm), the main treatment decision is based on renal anatomy and stone composition. She added that ureterorenoscopy is a reasonable option for small to mid-size stones or stones that have failed SWL.

Infertility

Lawrence Ross, Professor and Chairman of the Department of Urology at the University of Illinois, presented a number of papers on the management of infertility. In an informative lecture on the role of testis biopsy in the modern manage-

ment of infertility, Dr. Ross said there is little role for same. The differential of obstructive versus nonobstructive azoospermia can not be determined by physical examination of the testis and endocrine profiles. In a very interesting paper, Dr. Ross reviewed the cost/benefit analysis of the management of male infertility. He noted that in cases of severe male factors, such as azoospermia and severe oligospermia, the only treatment approach may be in vitro fertilization (IVF) or intracytoplasmic sperm injection (ICSI). But in the more common, treatable forms of male infertility, including varicocele, obstructive azoospermia, endocrine dysfunction, and infection, definitive therapy of these conditions may be more cost-effective. In a study by Neumann and associates,²⁴ charges per cycle for IVF from six facilities in the United States and estimated delivery rates from the published literature were evaluated. They observed that the cost per success for delivery with IVF was \$66,667 for the first cycle, increasing to \$114,286 by the sixth cycle. The cost was considerably greater in older women and if there was a male factor contributing to the

infertility. The Society for Assisted Reproductive Technology has reported that for the most recent reporting period (1998), when three embryos are transferred in an IVF cycle, the live birth rate is 37.7% for a woman 20 to 29 years of age, diminishing to 8.3% for women 40 to 44. The overall birth rate per cycle for all patients undergoing IVF remains at 25% and unfortunately has not changed over the last 5 years.

Schlegel²⁵ did a cost-effectiveness comparison of varicocele ligation to IVF/ICSI. The cost for varicolectomy was \$26,268 versus \$89,091, based on figures from Cornell University. They noted that the average published delivery rate for ICSI was 28% versus 30% for varicolectomy. Comparing similar results for vasectomy reversal, the cost for delivery was \$72,521 for sperm retrieval and ICSI versus \$25,475 for vasovasostomy. These data suggested that in cases of male-factor infertility procedures directed to the male generally

are more cost effective and allow natural conception.

Upcoming Meetings at Jackson Hole

The 2002 Jackson Hole Summer Meeting will be held July 27–August 2. Interested readers should contact Kay Christensen, Conference Coordinator, for more information at 307-856-5758.

References

1. Lui PD, Terris MK, McNeal JE, Stamey TA. Indications for ultrasound guided transition zone biopsies in the detection of prostate cancer. *J Urol.* 1995;153:1000–1003.
2. Liu JJ, Macy M, Lai Y, Terris MK. Critical evaluation of the current indications for transition zone biopsies. *Urology.* 2001;57:1117–1120.
3. Babaian RJ. Extended field prostate biopsy enhances cancer detection. *Urology.* 2000;55:453–456.
4. Bazinet M, Karakiewicz PI, Aprikian AG, et al. Value of systematic transition zone biopsies in the early detection of prostate cancer. *J Urol.* 1996;155:605–606.
5. Keetch DW, Catalona WJ. Prostatic transition zone biopsies in men with previous negative biopsies and persistently evaluated serum prostate specific antigen values. *J Urol.* 1995;154:1795–1797.
6. Morote J, Lopez M, Encabo G, de Torres I. Value of routine transition zone biopsies in patients undergoing ultrasound-guided sextant biopsies for the first time. *Eur Urol.* 1999;35:294–297.
7. Onder AU, Yalcin V, Arar O, et al. Impact of transition zone biopsies in detection and evaluation of prostate cancer. *Eur Urol.* 1998;33:542–548.
8. Iversen P, Tyrrell CJ, Kaisary AV, et al. Bicalutamide monotherapy compared with castration in patients with nonmetastatic locally advanced prostate cancer. 6.3 years of followup. *J Urol.* 2000;164:1579–1582.
9. Chodak G, Sharifi R, Kasimis B, et al. Single-agent therapy with bicalutamide: a comparison with medical or surgical castration in the treatment of advanced prostate carcinoma. *Urology.* 1995;46:849–855.
10. Bales GT, Chodak GW. A controlled trial of bicalutamide versus castration in patients with advanced prostate cancer. *Urology.* 1996;47:38–43.
11. Tyrrell CJ, Kaisary AV, Iversen P, et al. A randomized comparison of 'Casodex' (bicalutamide) 150 mg monotherapy versus castration in the treatment of metastatic and locally advanced prostate cancer. *Eur Urol.* 1998;33:447–456.
12. Boccardo F, Rubagotti A, Barichello M, et al. Bicalutamide monotherapy versus flutamide plus goserelin in prostate cancer patients: results of an Italian Prostate Cancer Project study. *J Clin Oncol.* 1999;17:2027–2038.
13. Nudell DM, Grossfeld GD, Weinberg VK, et al. Radiotherapy after radical prostatectomy: treatment outcomes and failure patterns. *Urology.* 1999;54:1049.
14. Grossfeld GD, Tigrani VS, Nudell D, et al. Management of a positive surgical margin after radical prostatectomy: decision analysis. *J Urol.* 2000;164:93–99;discussion 100.
15. Stein JP, Cote RJ, Freeman JA, et al. Indications for lower urinary tract reconstruction in women after cystectomy for bladder cancer: a pathologic

Main Points

- A prospective evaluation of 259 men with lower urinary tract symptoms (LUTS) who underwent transurethral resection of the prostate (TURP) demonstrated that a considerable portion of patients who have negative peripheral and transition zone biopsy may harbor significant carcinoma of the prostate at the time of TURP.
- A study comparing 33 men with isolated positive apical margins following radical prostatectomy to a cohort of controls concluded that, when controlling for other prognostic parameters, a positive margin at the apex is not associated with untoward prognosis.
- Univariate and multivariate analysis of the CaPSURE registry of patients in the United States showed that percentage of biopsies positive for cancer following radical prostatectomy is a significant predictor of both extracapsular disease and disease recurrence.
- A presentation on orthotopic urinary diversion in women requiring cystectomy noted that increase in experience has demonstrated this to be a safe and reliable method of urinary diversion in properly selected patients.
- After reviewing treatment options for ureteral strictures, one presenter concluded that endourologic management of benign ureteral and ureteroenteric strictures provides reasonable first-line therapy. Endoureterotomy has a higher success rate than balloon dilation, and the latter is exceedingly prone to failure for ureteroenteric strictures.
- A presentation regarding the outcome and management of lower pole calculi concluded that for small stones (<10 mm), shockwave lithotripsy is reasonable first-line therapy. For larger stones (>20 mm), percutaneous nephrolithotomy is preferred. For stones intermediate in size (11–20 mm), the main treatment decision is based on renal anatomy and stone composition.
- A review of cost/benefit analyses of the management of male infertility concluded that in cases of severe male factors, such as azoospermia and severe oligospermia, the only treatment approach may be in vitro fertilization or intracytoplasmic sperm injection; but in the more common, treatable forms of male infertility, including varicocele, obstructive azoospermia, endocrine dysfunction, and infection, definitive therapy of these conditions may be more cost-effective.

- ical review of female cystectomy specimens. *J Urol.* 1995;154:1329-1333.
16. Hubner WA, Irby P, Stoller ML. Natural history and current concepts for the treatment of small ureteral calculi. *Eur Urol.* 1993;24:172-176.
17. Leventhal EK, Rozanski TA, Crain TW, Deshon GE Jr. Indwelling ureteral stents as definitive therapy for distal ureteral calculi. *J Urol.* 1995;153:34-36.
18. Denstedt JD, Wollin TA, Sofer M, et al. A prospective randomized controlled trial comparing nonstented versus stented ureteroscopic lithotripsy. *J Urol.* 2001;165:1419-1422.
19. Deliveliotis C, Stavropoulos NI, Koutsokalis G, et al. Distal ureteral calculi: ureteroscopy vs. ESWL. A prospective analysis. *Int Urol Nephrol.* 1996;28:627-631.
20. Peschel R, Janetschek G, Bartsch G. Extracorporeal shock wave lithotripsy versus ureteroscopy for distal ureteral calculi: a prospective randomized study. *J Urol.* 1999;62:1909-1912.
21. Beckmann CF, Roth RA, Bihle W 3rd. Dilation of benign ureteral strictures. *Radiology.* 1989;172:437-441.
22. Chang R, Marshall FF, Mitchell S. Percutaneous management of benign ureteral strictures and fistulas. *J Urol.* 1987;137:1126-1131.
23. Lingeman JE, Siegel YI, Steele B, et al. Management of lower pole nephrolithiasis: a critical analysis. *J Urol.* 1994;151:663-667.
24. Neumann PJ, Gharib SD, Weinstein MC. The cost of a successful delivery with in vitro fertilization. *N Engl J Med.* 1994;4:239-243.
25. Schlegel PN. Is assisted reproduction the optimal treatment for varicocele-associated male infertility: a cost-effectiveness analysis. *Urology.* 1997;1:83-90.
26. Ornstein DK, Rao GS, Smith DS, Andriole GL. The impact of systematic prostate biopsy on prostate cancer incidence in men with symptomatic benign prostatic hyperplasia undergoing transurethral resection of the prostate. *J Urol.* 1997;157:880-884.
27. Blute ML, Gburek BM. Continent orthotopic urinary diversion in female patients: early Mayo Clinic experience. *Mayo Clin Proc.* 1998;73:501-507.
28. Cancrini A, De Carli P, Fattahi H, et al. Orthotopic ileal neobladder in female patients after radical cystectomy: 2-year experience. *J Urol.* 1995;153:956-958.
29. Hautmann RE, Paiss T, de Petriconi R. The ileal neobladder in women: 9 years of experience with 19 patients. *J Urol.* 1996;155:76-81.
30. Jarolim L, Babjuk M, Hanus T, et al. Female urethra-sparing cystectomy and orthotopic bladder replacement. *Eur Urol.* 1997;31:173-177.
31. Mills RD, Studer UE. Female orthotopic bladder substitution: a good operation in the right circumstances. *J Urol.* 2000;163:1501-1504.
32. Shimogaki H, Okada H, Fujisawa M, et al. Long-term experience with orthotopic reconstruction of the lower urinary tract in women. *J Urol.* 1999;161:573-577.
33. Stenzl A, Colleselli K, Bartsch G. Update of urethra-sparing approaches in cystectomy in women. *World J Urol.* 1997;15:134-138.
34. Johnson CD, Oke EJ, Dunnick NR, et al. Percutaneous balloon dilation of ureteral strictures. *AJR Am J Roentgenol.* 1987;148:181-184.
35. Smith AD. Management of iatrogenic ureteral strictures after urological procedures. *J Urol.* 1988;140:1372-1374.
36. O'Brien WM, Maxted WC, Pahira JJ. Ureteral stricture: experience with 31 cases. *J Urol.* 1988;140:737-740.
37. Beckmann CF, Roth RA. Secondary ureteropelvic junction stricture: percutaneous dilation. *Radiology.* 1987;164:365-367.
38. Kramolowsky EV, Tucker RD, Nelson CM. Management of benign ureteral structures: open surgical repair or endoscopic dilation? *J Urol.* 1989;141:285-286.
39. Netto NR Jr, Ferreira U, Lemos GC, Claro JF. Endourological management of ureteral strictures. *J Urol.* 1990;144:631-634.
40. Ravery V, de la Taille A, Hoffmann P, et al. Balloon catheter dilation in the treatment of ureteral and ureteroenteric stricture. *J Endourol.* 1998;12:335-340.
41. Schneider AW, Conrad S, Busch R, Otto U. The cold-knife technique for endourological management of stenoses in the upper urinary tract. *J Urol.* 1991;146:961-965.
42. Meretyk S, Albala DM, Clayman RV, et al. Endouretotomy for treatment of ureteral strictures. *J Urol.* 1992;147:1502-1506.
43. Yamada S, Ono Y, Ohsima S, Miyake K. Transurethral ureteroscopic ureterotomy assisted by a prior balloon dilation for relieving ureteral strictures. *J Urol.* 1995;153:1418-1421.
44. Cornud F, Lefebvre JF, Chretien Y, et al. Percutaneous transrenal electro-incision of uretero-intestinal anastomotic strictures: long-term results and comparison of fluoroscopic and endoscopic guidance. *J Urol.* 1996;155:1575-1578.
45. Preminger GM, Clayman RV, Nakada SY, et al. A multicenter clinical trial investigating the use of a fluoroscopically controlled cutting balloon catheter for the management of ureteral and ureteropelvic junction obstruction. *J Urol.* 1997;157:1625-1629.
46. Singal RK, Denstedt JD, Razvi HA, Chun SS. Holmium: YAG laser endouretotomy for treatment of ureteral stricture. *Urology.* 1997;50:875-880.
47. Wolf JS Jr, Elashry OM, Clayman RV. Long-term results of endouretotomy for benign ureteral and ureteroenteric strictures. *J Urol.* 1997;158:759-764.